6. Christiania omegns Phanerogamer og Bregner med Angivelse af deres udbredelse samt en indledning om vegetationens afhængighed af underlaget af A. Blytt.

From Prof. Owen:-

On a new species of Sthenurus.

PAPERS READ.

Description of a new species of Ptilotis, from Torres Straits. By E. P. Ramsay, F.L.S.

PTILOTIS GERMANA. sp. nov.

Adult male. The front and sides of the head, lores, throat and chest, and the margins of the shoulders on the under side, yellow; ear-coverts yellow, with a narrow stripe of black, commencing almost at the angle of the mouth, passing through them, and below which they are of a brighter yellow; under wing-coverts pale buff; chest, breast, and remainder of the under surface pale yellow, indistinctly striped down the centre with dull brown, centre of the abdomen and under tail-coverts citron yellow, the greater series of the under tail-coverts striped down the centre with brown; back of the head, hind neck, and all the upper surface of the body dull brown, with a faint tinge of olive on the rump; wings and tail dark brown all the feathers broadly margined on the outer webs with bright olive yellow; bill black, legs brown.

Total length, 5.5; wings, 3.1; tail, 2.5; tarsi, 0.8; bill from forehead, 0.7; from gape, 0.75.

Hab. Torres Straits.

This species comes close to *P. ornatus* (Gould), on the one hand from the stripes of the chest, but has not the blackish lores of that species; on the other hand it resembles *P. flavescens* of the same author, but may be distinguished from it by the dull brown of the back, and stripes on the under surface.